

Right Eye / Left Eye

Cataract Surgery Post-Operative Eye Drop Schedule

	1 st Week	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
1 x a Day	Prolensa	0	0	0	0	0	0	0
4 x a Day	Durezol	0000	0000	0000	0000	0000	0000	0000
3 x a Day	Besivance	000	000	000	000	000	000	000
	2 nd Week	Day 8	Day 9	Day 10	Day 11	Day 12	Day 13	Day 14
1 x a Day	Prolensa	0	0	O	0	O	O	0
2 x a Day	Durezol	0 0	0 0	0 0	0 0	0 0	0 0	0 0
	3 rd Week	Day 15	Day 16	Day 17	Day 18	Day 19	Day 20	Day 21
1 x a Day	Prolensa	O O	O O	O 0	O O	O O	O Day 20	O Day 21
2 x a Day	Durezol	0 0	0 0	0 0	0 0	0 0	0 0	0 0
	4th Mark	Day 22	Day 22	Day 24	Day 25	Day 26	Day 27	Day 20
1 x a Day	4 th Week Prolensa	Day 22 O	Day 23	Day 24 O	Day 25	Day 26 O	Day 27 O	Day 28 O
2 x a Day	Durezol	0 0	0 0	0 0	0 0	0 0	0 0	0 0
						_		
1 x a Day	5 th Week	Day 29	Day 30	Day 31	Day 32	Day 33	Day 34	Day 35
	Prolensa	0	0	0	0	0	0	0
2 x a Day	Durezol	0 0	0 0	0 0	0 0	0 0	0 0	0 0
	6 th Week	Day 36	Day 37	Day 38	Day 39	Day 40	Day 41	Day 42
1 x a Day	Prolensa	0	0	0	0	0	0	0
2 x a Day	Durezol	0 0	0 0	0 0	0 0	0 0	0 0	0 0
	7 th Week	Day 43	Day 44	Day 45	Day 46	Day 47	Day 48	Day 49
1 x a Day	Prolensa	O	0	O	0	O	0	O
2 x a Day	Durezol	0 0	0 0	0 0	0 0	0 0	0 0	0 0
	8 th Week	Day 50	Day 54	Day 52	Dov.52	Day-54	Dov-FF	Dov-FC -
1 x a Day	Prolensa	Day 50 O	Day 51	Day 52 O	Day 53	Day 54 O	Day 55 O	Day 56
2 x a Day	Durezol	0 0	0 0	0 0	0 0	0 0	0 0	0 0
	SYSTANE /	RETAINE	Use as much or as little as you need for comfort from irritation.					

Please call our office if you experience sudden vision loss or pain.



- Wait 3-5 minutes in between each eye drop
- You may sleep on your operated side. Wear your protective shield for the first week when you nap and sleep.
- Do not rub your operated eye.
- You may resume most of your NORMAL activities.
- Bending is ok.
- Limit any activity which causes excessive strain or repeated jarring for one week.
- You may wash your face and hair. Be careful not to get shampoo in your operated eye.
- No swimming for 4 weeks.
- Wear your dark plastic glasses as desired. They offer good protection and decrease glare.

Please call our office at 713-791-9494 if you experience sudden vision loss or pain.



Cataract Pre-Operative Instructions

If you are currently on blood thinners or aspirin please DO NOT discontinue prior to surgery.

		3 DAYS	3 DAYS PRIOR TO SURGERY			
		DAY 1	DAY 2	DAY 3		
1 x a Day	Prolensa (GREY CAP)	O am	O am	O am		
2 x a Day	Durezol (PINK CAP)	O O am pm	O O am pm	O O am pm		
3 x a Day	Besivance (BEIGE CAP)	O O O am noon pm	O O O am noon pm	O O O am noon pm		

MORNING OF SURGERY					
	Prolensa (GREY CAP)	O am			
	Durezol (PINK CAP)	O am			
	Besivance (BEIGE CAP)	O am			

- ♦ Wait 3-5 minutes in between each eye drop.
- ♦ Save these medications. You will need them after the surgery.
- ◆ DO NOT EAT OR DRINK ANYTHING AFTER MIDNIGHT THE DAY BEFORE SURGERY.
- ◆ Take all heart and/or blood pressure medications that you would normally take on the morning of surgery with a small sip of water.
- Wear comfortable clothes (short sleeves and front buttons if possible).
- ◆ BRING YOUR PICTURE ID AND INSURANCE CARDS ON THE DAY OF SURGERY.
- ♦ You are REQUIRED to return to our office the day after surgery.