

**CATARACT COUMPOUNDED MEDICATION INSTRUCTIONS 13**  
**BEFORE AND AFTER SURGERY**

**Before Surgery**

**Pred-Nepafenac** → Instill 1 drop three times a day in the surgical eye(s) beginning the day before your procedure.

**After Surgery**

**Pred-Nepafenac** → Instill 1 drop three times a day in the surgical eye(s) for one week, then one drop two times a day until the bottle is completely empty.

PLEASE USE PRESERVATIVE FREE ARTIFICIAL TEARS AT LEAST EVERY HOUR WHILE AWAKE FOR THE FIRST 2-3 DAYS AND AS NEEDED FOR DISCOMFORT.

Istent/CYPASS glaucoma patients: Continue ALL of your glaucoma drops as you have been doing. You will continue to use them throughout your post op care. Your surgeon will not make any changes to your glaucoma medication routine until he feels your eye(s) are healed from cataract surgery to safely do so.

**IMPORTANT: PLEASE WEAR SHIELD AT BEDTIME FOR ONE WEEK.**

If you have any questions, please feel free to contact us at **713-580-2500**

For after hours **EMERGENCIES**, you may contact your surgeon directly through our emergency hotline:

**832-416-7340**  
**Dr. Jennifer Melton**